arXiv:math/0701725v3 [math.GT] 25 Oct 2008

ENDING LAMINATIONS AND CANNON-THURSTON MAPS

MAHAN MJ

Abstract. In earlier work, we had shown that Cannon-Thurston maps exist for Kleinian surface groups. In this paper we prove that pre-images of points are precisely end-points of leaves of the ending lamination whenever the Cannon-Thurston map is not one-to-one. In particular, the Cannon-Thurston map is finite-to-one. This completes the proof of the conjectural picture of Cannon-Thurston maps.

AMS Subject Classification: 57M50

Contents

1. Introduction and Preliminaries 1
1.1. Statement of Results 1
1.2. Ideal points are identified by Cannon-Thurston Maps 2
1.3. Hyperbolic Metric Spaces 4
1.4. Geodesic Laminations 5
2. Split Geometry 5
2.1. Hyperbolic Ladders and Retractions 8
3. Laminations 10
3.1. Leaves of Laminations 10
3.2. Rays Coarsely Separate Ladders 12
3.3. Main Theorem 14
References 15

1. Introduction and Preliminaries

1.1. Statement of Results.

In earlier work ([11] [13]) we showed:

Theorem 1.1. [11] [13] *Let* $\rho : \pi_1(S) \rightarrow PSL_2(C)$ *be a discrete faithful representation of a surface group with or without punctures, and without accidental parabolics. Let* $M = \mathbb{H}^3/\rho(\pi_1(S))$*. Let* i *be an embedding of* S *in* M *that induces a homotopy equivalence. Then the embedding* $\tilde{i} : \widetilde{S} \rightarrow \widetilde{M} = \mathbb{H}^3$ *extends continuously to a map* $\hat{i} : \mathbb{D}^2 \rightarrow \mathbb{D}^3$*. Further, the limit set of* $\rho(\pi_1(S))$ *is locally connected.*

This generalizes the first part of the next theorem due to Cannon and Thurston [3] (for 3 manifolds fibering over the circle) and Minsky [9] (for bounded geometry closed surface Kleinian groups):

Theorem 1.2. Cannon-Thurston [3], **Minsky** [9] *Suppose a closed surface group* $\pi_1(S)$ *of bounded geometry acts freely and properly discontinuously on* $\mathbb{H}^3$

by isometries. Then the inclusion $\widetilde{i} : \widetilde{S} \to \mathbb{H}^3$ *extends continuously to the boundary. Further, pre-images of points on the boundary are precisely ideal boundary points of a leaf of the ending lamination, or ideal boundary points of a complementary ideal polygon whenever the Cannon-Thurston map is not one-to-one.*

In this paper, we generalize the second part of the above theorem to arbitrary Kleinian closed surface groups.

Main Theorem 1.3: *Suppose a closed surface group* $\pi_1(S)$ *acts freely and properly discontinuously on* $\mathbb{H}^3$ *by isometries. Then the inclusion* $\widetilde{i} : \widetilde{S} \to \mathbb{H}^3$ *extends continuously to the boundary. Further, pre-images of points on the boundary are precisely ideal boundary points of a leaf of the ending lamination, or ideal boundary points of a complementary ideal polygon whenever the Cannon-Thurston map is not one-to-one.*

In passing from Theorem 1.2 to Theorem 1.3 we have removed the hypothesis of *bounded geometry.*

Acknowledgements: The author would like to thank an anonymous referee of an earlier version of this paper for pointing out some errors and suggesting corrections.

1.2. **Ideal points are identified by Cannon-Thurston Maps.**

We would like to know exactly which points are identified by the Cannon-Thurston map, whose existence is assured by Theorem 1.1. Let ∂i denote restriction to the boundary S^1_∞ of $\hat{i}$, the continuous extension to the boundary in Theorem 1.1.

We shall prove the easy direction of the following

Theorem 1.3. *Let* $\partial i(a) = \partial i(b)$ *for* $a, b \in S^1_\infty$ *be two distinct points that are identified by the Cannon-Thurston map corresponding to a simply degenerate closed surface group (without accidental parabolics). Then* a, b *are either ideal end-points of a leaf of a lamination, or ideal boundary points of a complementary ideal polygon. Conversely, if* a, b *are either ideal end-points of a leaf of a lamination, or ideal boundary points of a complementary ideal polygon, then* $\partial i(a) = \partial i(b)$.

It is easy to show the converse direction in Theorem 1.3 above. (cf. Lemma 3.5 of [10]):

Proposition 1.4. *Let* u, v *be either ideal end-points of a leaf of a lamination, or ideal boundary points of a complementary ideal polygon. Then* $\partial i(u) = \partial i(v)$.

Proof: (cf. Lemma 3.5 of [10]. See also [13])
Take a sequence of short geodesics s_i exiting the end. Let a_i be geodesics in the intrinsic metric on the convex core boundary S (of M) freely homotopic to s_i. (We use the hypothesis of simple degeneracy very mildly here to fix a base pleated surface.) We assume further that a_i's are simple. Join a_i to s_i by the shortest geodesic t_i in M connecting the two curves. Then the collection a_i converges to the ending lamination on S. Also, in the universal cover, we obtain segments $a_{fi} \subset \widetilde{S}$ which are finite segments whose end-points are identified by the covering map $P : \widetilde{M} \to M$. We also assume that P is injective restricted to the interior of a_{fi}'s mapping to a_i. Similarly there exist segments $s_{fi} \subset \widetilde{M}$ which are finite segments whose end-points are identified by the covering map $P : \widetilde{M} \to M$. We also assume that P is injective restricted to the interior of a_{fi}'s. The finite segments

s_{fi} and a_{fi} are chosen in such a way that there exist lifts t_{1i}, t_{2i}, joining end-points of a_{fi} to corresponding end-points of s_{fi}. The union of these four pieces looks like a trapezium (See figure below, where we omit subscripts for convenience).



Figure 1: *Trapezium*

Next, given any leaf λ of the ending lamination, we may choose translates of the finite segments a_{fi} (under the action of $\pi_1(S)$) appropriately, such that a_{fi} converge to λ in (the Hausdorff/Chabauty topology on closed subsets of) $\mathbf{H}^2$. For each a_{fi}, let

$$\beta_{fi} = t_{1i} \circ s_{fi} \circ \overline{t_{2i}}$$

where $\overline{t_{2i}}$ denotes t_{2i} with orientation reversed. Then β_{fi}'s are uniform hyperbolic quasigeodesics in $\widetilde{M}$. If the translates of a_{fi} we are considering have end-points lying outside large balls around a fixed reference point $p \in \widetilde{S}$, it is easy to check that β_{fi}'s lie outside large balls about p in $\widetilde{M}$.

At this stage we invoke the existence theorem for Cannon-Thurston maps, Theorem 1.1. Since a_{fi}'s converge to λ and there exist uniform hyperbolic quasigeodesics β_{fi}, joining the end-points of a_{fi} and exiting all compact sets, it follows that $\partial i(u) = \partial i(v)$, where a, b denote the boundary points of λ. The Proposition follows from

1) the fact (due to Bonahon [1]) that surface groups are tame and that for M there exist simple closed curves on S whose geodesic realizations exit the end.

2) the fact (due to Thurston) that any sequence of simple closed curves on S with geodesic realizations exiting the end, converges in S to the ending lamination ([16] Ch. 9)

3) consequently, any leaf of the ending lamination may be realized as the limit of geodesic segments a_{fi}

4) If we define u, v to be equivalent if they are the end-points of a leaf of the ending lamination, then the transitive closure of this relation has as elements of an equivalence class

a) either ideal end-points of a leaf of a lamination,

b) or ideal boundary points of a complementary ideal polygon,

c) or a single point in S^1_∞ which is not an end-point of a leaf of a lamination. $\square$

Remark: Proposition 1.4 shows that the existence of a Cannon-Thurston map automatically guarantees that end-points of leaves of the ending lamination are identified by the Cannon-Thurston map.

Definition 1.5. *Let H be a finitely presented group acting on a hyperbolic space X with quotient M. Let X_H be a 2-complex with fundamental group H, and $i : X_H \to M$ be a map inducing an isomorphism of fundamental groups. Then i lifts*

to $\tilde{i} : \tilde{X_H} \to X$*. A bi-infinite geodesic* λ *in* $\tilde{X_H} \subset X$ *will be called a* **leaf of the abstract ending lamination** *for* $i : X_H \to M$*, if*
1) there exists a set of geodesics σ_i *in* M *exiting every compact set*
2) there exists a set of geodesics α_i *in* X_H *with* $i(\alpha_i)$ *freely homotopic to* σ_i
3) there exist finite lifts $\widetilde{\alpha_i}$ *of* α_i *in* $(\widetilde{X_H})$ *such that the natural covering map* $\Pi : \widetilde{X_H} \to X_H$ *is injective away from end-points of* $\widetilde{\alpha_i}$
4) $\widetilde{\alpha_i}$ *converges to* λ *in the Chabauty topology*

Proposition 1.4 and its proof readily generalize to

Proposition 1.6. *Suppose* H *is hyperbolic and* $\tilde{i} : \widetilde{X_H} \to X$ *extends to a Cannon-Thurston map on boundaries. Let* u, v *be end-points of a leaf of an abstract ending lamination. Then* $\partial i(u) = \partial i(v)$*.*

1.3. **Hyperbolic Metric Spaces.**

Let (X, d_X) be a hyperbolic metric space and Y be a subspace that is hyperbolic with the inherited path metric d_Y. By adjoining the Gromov boundaries ∂X and ∂Y to X and Y, one obtains their compactifications $\widehat{X}$ and $\widehat{Y}$ respectively.

Let $i : Y \to X$ denote inclusion.

Definition 1.7. *Let* X *and* Y *be hyperbolic metric spaces and* $i : Y \to X$ *be an embedding. A* **Cannon-Thurston map** $\hat{i}$ *from* $\widehat{Y}$ *to* $\widehat{X}$ *is a continuous extension of* i*.*

The following Lemma is a 'quasification' of the fact that geodesics converging to the same point on $\partial\Gamma_H$ are asymptotic [See [7], pg. 117].

Relative Hyperbolicity and Electric Geometry

We refer the reader to Farb [5] for terminology and details on relative hyperbolicity and electric geometry.

Let X be a δ-hyperbolic metric space, and $\mathcal{H}$ a family of C-quasiconvex, D-separated, collection of subsets. Then by work of Farb [5], X_{el} obtained by electrocuting the subsets in $\mathcal{H}$ is a $\Delta = \Delta(\delta, C, D)$ -hyperbolic metric space. (Recall [5] that **electrocution** means constructing an auxiliary space $X_{el} = X \bigcup_{H \in \mathcal{H}} (H \times I)$ with $H \times \{0\}$ identified to $H \subset X$ and $H \times \{1\}$ equipped with the zero metric. This is a geometric 'coning' construction.) Now, let $\alpha = [a, b]$ be a hyperbolic geodesic in X and β be an electric P-quasigeodesic without backtracking joining a, b. Starting from the left of β, replace each maximal subsegment, (with end-points p, q, say) lying within some $H \in \mathcal{H}$ by a hyperbolic geodesic $[p, q]$. The resulting **connected** path β_q is called an *electro-ambient representative* in X.

In [12] we noted that β_q *need not be a hyperbolic quasigeodesic.* However, we did adapt Proposition 4.3 of Klarreich [8] to obtain the following:

Lemma 1.8. *(See Proposition 4.3 of* [8]*, also see Lemma 3.10 of* [12]*) Given* δ*,* C, D, P *there exists* C_3 *such that the following holds:*
Let (X, d) *be a* δ*-hyperbolic metric space and* $\mathcal{H}$ *a family of* C*-quasiconvex,* D*-separated collection of quasiconvex subsets. Let* (X, d_e) *denote the electric space obtained by electrocuting elements of* $\mathcal{H}$*. Then, if* α, β_q *denote respectively a hyperbolic geodesic and an electro-ambient* P*-quasigeodesic with the same end-points, then* α *lies in a (hyperbolic)* C_3 *neighborhood of* β_q*.*

1.4. **Geodesic Laminations.** For a discussion of geodesic laminations (or simply laminations as we shall call them), we refer the reader to [14], [2], [16], [4]. For a discussion on dual $\mathbb{R}$-trees, see [15].

The space of filling laminations which we denote $\mathcal{FL}$ are the measure classes of measured laminations Λ for which all complementary regions of the support $|\Lambda|$ are simply connected. The quotient of $\mathcal{FL}$ by forgetting the measures will be denoted $\mathcal{EL}$ and is the space of *ending laminations*. For the purposes of this paper, the only laminations we shall be interested in are those without simple closed leaves, i.e. *arational laminations*. We shall therefore assume that all filling laminations and ending laminations are arational. A useful fact is that an arational laminations is that they are minimal, i.e. the closure (in the Hausdorff topology) of **any** of its leaves is the whole lamination. A useful fact about minimal laminations is the following. We can identify a minimal lamination Λ with a closed invariant (under $\pi_1(S)$) subset of the set of *unordered pairs* in $(S^1_\infty \times S^1_\infty \setminus \Delta)/\mathbf{R_a}$, where Δ denotes the diagonal $\mathbf{R_a}$ is the relation identifying (a, b) with (b, a).

Lemma 1.9. *Let Λ be a minimal geodesic lamination on a surface S. Let I be an embedded (closed) interval in S transverse to Λ. Let $\widetilde{\Lambda}$ denote the union of all lifts of leaves of Λ to the universal cover $\widetilde{S}$. Let $\widetilde{I}$ denote the union of all lifts of I to $\widetilde{S}$. Define two leaves of $\widetilde{\Lambda}$ to be equivalent if both of them intersect a single element of $\widetilde{I}$. Then the limit set of any connected component of the transitive closure of this relation contains a pole.*

Proof: Let $\mathcal{T}$ be the $\mathbb{R}$-tree dual to $\widetilde{\Lambda}$. Then $I \subset \widetilde{S}$ projects to an embedded non-trivial interval in $\mathcal{T}$ under the quotient map q that identifies leaves of $\widetilde{\Lambda}$ to points. Then the orbit of I under $\pi_1(S)$ acting on $\mathcal{T}$ is a forest, in fact a sub-forest $\mathcal{F}$ of $\mathcal{T}$. Look at the connected component $\mathcal{T}_1$ of $\mathcal{F}$ containing I.

If $g\mathcal{T}_1 \cap \mathcal{T}_1 \neq \emptyset$, then $g\mathcal{T}_1 \subset \mathcal{T}_1$. Hence $g^{-1}\mathcal{T}_1 \cap \mathcal{T}_1 \neq \emptyset$, and we finally have that $\mathcal{T}_1$ is invariant under g^n for all integers n. This shows that $q^{-1}(\mathcal{T}_1) \subset \widetilde{S}$ contains the pole corresponding to the infinite order element g.

Thus we need finally the existence of a g as in the previous paragraph. It suffices to show that for any non-trivial I, there exists $g \in \pi_1(S)$ such that $gI \cap I \neq \emptyset$. But this follows from minimality of Λ, using the fact that each leaf is dense in Λ, and hence that there exists $g \in \pi_1(S)$ such that gI and I are transverse to a *common leaf* λ of $\widetilde{\Lambda}$. $\Box$

2. Split Geometry

We shall briefly recall the construction of models of *split geometry* from [11]. We shall also need the construction of certain quasiconvex ladder-like sets B_λ.

Topologically, a **split subsurface** S^s of a surface S is a (possibly disconnected, proper) subsurface with boundary such that $S - S^s$ consists of a non-empty family of non-homotopic essential annuli, which in turn are not homotopic into the boundary of S^s.

Geometrically, we assume that S is given some finite volume hyperbolic structure. A split subsurface S^s of S has bounded geometry, i.e.
1) each boundary component of S^s is of length ϵ_0, and is in fact a component of the boundary of $N_k(\gamma)$, where γ is a hyperbolic geodesic on S, and $N_k(\gamma)$ denotes its k-neighborhood.

2) For any closed geodesic β on S, either $\beta \subset S - S^s$, or, the length of any component of $\beta \cap (S - S^s)$ is greater than ϵ_0.

Topologically, a **split block** $B^s \subset B = S \times I$ is a topological product $S^s \times I$ for some *connected* S^s. However, its upper and lower boundaries need not be $S^s \times 1$ and $S^s \times 0$. We only require that the upper and lower boundaries be *split subsurfaces* of S^s. This is to allow for Margulis tubes starting (or ending) within the split block. Such tubes would split one of the horizontal boundaries but not both. We shall call such tubes **hanging tubes**. Connected components of split blocks are called **split components**. We demand that there is a *non-empty* collection of Margulis tubes splitting a split block.

If a block has no Margulis tubes splitting it and the block has no hanging tubes, it is called a **thick block**.

We further require that the distance between horizontal boundary components is at least 1, i.e. for a component R of S^s $d(R \times 0, R \times 1) \geq 1$. We define the **thickness** of a split block to be the supremum of the lengths of $x \times I$ for $x \in S^s$ and demand that it be finite (which holds under all reasonable conditions, e.g. a smooth metric).

See figure below, where the left split component has four hanging tubes and the right split component has two hanging tubes. The vertical space between the components is the place where a Margulis tube splits the split block into two split components.



Figure 2: *Split Components of Split Block with hanging tubes*

Geometrically, we require that the metric on a split block induces a path metric on its *upper and lower horizontal* boundary components, which are subsurfaces of $S^s \times \partial I$, such that each horizontal boundary component is a (geometric) split surface. Further, the metric on B^s induces on each *vertical* boundary component of a Margulis tube $\partial S^s \times I$ the product metric. Each boundary component for Margulis tubes that 'travel all the way from the lower to the upper boundary' is an annulus of height equal to length of I. We demand further that *hanging tubes* have *length uniformly bounded below by* $\eta_0 > 0$. Further, each such annulus has cross section a round circle of length ϵ_0. This leaves us to decide the metric on lower and upper boundaries of hanging tubes. Such boundaries are declared to have a metric equal to that on $S^1 \times [-\eta, \eta]$, where S^1 is a round circle of length ϵ_0 and η is a sufficiently small number.

As noted in the definition of *split components*, we *do not* require that the upper (or lower) horizontal boundary of a split component K be connected. This happens due to the presence of *hanging tubes*.

We further require that the 'vertical boundaries' (corresponding to Euclidean annuli) of split components be uniformly (independent of choice of a block and a split component) quasiconvex in the corresponding split component. Note that the boundary of each split block has an intrinsic metric that is flat and corresponds to a Euclidean torus. We define a **welded split block** to be a split block with identifications as follows: Components of $\partial S^s \times 0$ are glued together if and only if they correspond to the same geodesic in $S - S^s$. The same is done for components of $\partial S^s \times 1$. A simple closed curve that results from such an identification shall be called a **weld curve**. For hanging tubes, we also weld the boundary circles of their *lower or upper boundaries* by simply collapsing $S^1 \times [-\eta, \eta]$ to $S^1 \times \{0\}$.

Let the metric product $S^1 \times [0,1]$ be called the **standard annulus** if each horizontal S^1 has length ϵ_0. For hanging tubes the standard annulus will be taken to be $S^1 \times [0, 1/2]$.

Next, we require another pseudometric on B which we shall term the **tube-electrocuted metric**. We first define a map from each boundary annulus $S^1 \times I$ (or $S^1 \times [0, 1/2]$ for hanging annuli) to the corresponding standard annulus that is affine on the second factor and an isometry on the first. Now glue the mapping cylinder of this map to the boundary component. The resulting 'split block' has a number of standard annuli as its boundary components. Call the split block B^s with the above mapping cylinders attached, the *stabilized split block* B^{st}.

Glue boundary components of B^{st} corresponding to the same geodesic together to get the **tube electrocuted metric** on B as follows. Suppose that two boundary components of B^{st} correspond to the same geodesic γ. In this case, these boundary components are both of the form $S^1 \times I$ or $S^1 \times [0, \frac{1}{2}]$ where there is a projection onto the horizontal S^1 factor corresponding to γ. Let $S^1_l \times J$ and $S^1_r \times J$ denote these two boundary components (where J denotes I or $[0, \frac{1}{2}]$). Then each $S^1 \times \{x\}$ has length ϵ_0. Glue $S^1_l \times J$ to $S^1_r \times J$ by the natural 'identity map'. Finally, on each resulting $S^1 \times \{x\}$ put the zero metric. Thus the annulus $S^1 \times J$ obtained via this identification has the zero metric in the *horizontal direction* $S^1 \times \{x\}$ and the Euclidean metric in the *vertical direction* J. The resulting block will be called the **tube-electrocuted block** B_{tel} and the pseudometric on it will be denoted as d_{tel}. Note that B_{tel} is homeomorphic to $S \times I$. The operation of obtaining a *tube electrocuted block and metric* (B_{tel}, d_{tel}) from a split block B^s shall be called *tube electrocution.* Note that a tube electrocuted block is homeomorphic to $S \times I$.

A lift of a split component to the universal cover of the block $B = S \times I$ or $B_{tel} = S \times I$ shall be termed a **split component** of $\tilde{B}$ or $\tilde{B_{tel}}$.

Also, let d_G be the (pseudo)-metric obtained by electrocuting the collection $\widetilde{K}$ in $\tilde{B_{tel}}$. d_G will be called the the **graph metric**.

DEFINITION-THEOREM (SPLIT GEOMETRY)

We constructed in [13] the following from the Minsky model for a simply or totally degenerate surface group:

1) A sequence of split surfaces S^s_i exiting the end(s) of M.

2) A collection of Margulis tubes $\mathcal{T}$.

3) For each complementary annulus of S^s_i with core σ, there is a Margulis tube T whose core is freely homotopic to σ and such that T intersects the level i. (What this roughly means is that there is a T that contains the complementary annulus.)

4) For all i, either there exists a Margulis tube splitting both S^s_i and S^s_{i+1} and hence B^s_i, or else B_i is a thick block.

5) $T \cap S_i^s$ is either empty or consists of a pair of boundary components of S_i^s that are parallel in S_i.
6) There is a uniform upper bound n on the number of surfaces that T splits.
7) Each split component $\widetilde{K} \subset \widetilde{B_i} \subset \widetilde{M}$ is (not necessarily uniformly) quasiconvex in the hyperbolic metric on $\widetilde{M}$.
8) Equip $\widetilde{M}$ with the *graph-metric* d_G obtained by electrocuting each split component $\widetilde{K}$. Then the convex hull $CH(\widetilde{K})$ of any split component $\widetilde{K}$ has uniformly bounded diameter in the metric d_G. We say that the components $\widetilde{K}$ are uniformly graph-quasiconvex. It follows that $(\widetilde{M}, d_G)$ is a hyperbolic metric space.

Definition 2.1. *A model manifold satisfying conditions (1)-(8) above is said to have* **split geometry**.

2.1. Hyperbolic Ladders and Retractions. Admissible Paths

Admissible paths consist of the following:
1) Horizontal segments along some $\widetilde{S} \times \{i\}$ $i = \{0, 1\}$.
2) Vertical segments $x \times [0, 1]$.

Ladders in Building Blocks

We shall need to construct a set B_λ containing λ and a retraction Π_λ of $\widetilde{M}$ onto it. Π_λ will have the property that it does not stretch distances much. This will show that B_λ is quasi-isometrically embedded in $\widetilde{M}$.

To begin, we describe the construction of B_λ restricted to a building block B.

Construction of $B_\lambda(B)$ - Thick Block

Let the thick block be the universal curve over a Teichmuller geodesic $[\alpha, \beta]$. Let S_α denote the hyperbolic surface over α and S_β denote the hyperbolic surface over β.

First, let $\lambda = [a, b]$ be a geodesic segment in $\widetilde{S}$. Let λ_{B0} denote $\lambda \times \{0\}$.

Next, let ψ be the lift of the 'identity' map from $\widetilde{S_\alpha}$ to $\widetilde{S_\beta}$. Let Ψ denote the induced map on geodesics and let $\Psi(\lambda)$ denote the hyperbolic geodesic joining $\psi(a), \psi(b)$. Let λ_{B1} denote $\Psi(\lambda) \times \{1\}$.

For the universal cover $\widetilde{B}$ of the thick block B, define $B_\lambda(B) = \bigcup_{i=0,1} \lambda_{Bi}$.

Definition: Each $\widetilde{S} \times i$ for $i = 0, 1$ will be called a **horizontal sheet** of $\widetilde{B}$ when B is a thick block.

Construction of $B_\lambda(B)$ - Split Block

First, recall that $\lambda = [a, b]$ is a geodesic segment in $\widetilde{S}$. Let λ_{B0} denote $\lambda \times \{0\}$.

Next, let λ_{Gel} denote the electric geodesic joining a, b in the electric pseudometric on $\widetilde{S}$ obtained by electrocuting lifts of split components. Let λ_{B1} denote $\lambda_{Gel} \times \{0\}$.

Third, recall that $\tilde{\phi}$ is the lift of a component preserving diffeomorphism ϕ to $\widetilde{S}$ equipped with the electric metric d_{Gel}. Let $\tilde{\Phi}$ denote the induced map on geodesics, i.e. if $\mu = [x, y] \subset (\widetilde{S}, d_{Gel})$, then $\tilde{\Phi}(\mu) = [\phi(x), \phi(y)]$ is the geodesic joining $\phi(x), \phi(y)$. Let λ_{B2} denote $\tilde{\Phi}(\lambda_{Gel}) \times \{1\}$.

Fourthly, let $\tilde{\Phi}(\lambda)$ denote the hyperbolic geodesic joining $\phi(a), \phi(b)$. Let λ_{B3} denote $\tilde{\Phi}(\lambda) \times \{1\}$.

For the universal cover $\widetilde{B}$ of the thin block B, define $B_\lambda(B) = \bigcup_{i=0,\cdots,3} \lambda_{Bi}$.

Definition: Each $\widetilde{S} \times i$ for $i = 0 \cdots 3$ will be called a **horizontal sheet** of $\widetilde{B}$ when B is a thick block.

Modifications for Bi-infinite Geodesics
Only a few minor modifications are necessary for the above construction to go through for bi-infinite geodesics λ. In both a thick block and a split block, the maps ψ and ϕ are quasi-isometries (not necessarily uniform for split blocks). This induces maps Ψ and Φ on geodesics. If λ is an infinite geodesic (semi-infinite or bi-infinite), its image is a quasigeodesic and hence lies in a bounded neighborhood of a geodesic, which we denote as $\Psi(\lambda)$ or $\Phi(\lambda)$. (This is well-defined up to asymptote classes, i.e. one geodesic from an asymptote class is chosen.) With this understanding, the construction of $B_\lambda(B)$ goes through *mutatis mutandis.*

Construction of $\Pi_{\lambda,B}$ - Thick Block
On $\widetilde{S} \times \{0\}$, let Π_{B0} denote nearest point projection onto λ_{B0} in the path metric on $\widetilde{S} \times \{0\}$.

On $\widetilde{S} \times \{1\}$, let Π_{B1} denote nearest point projection onto λ_{B1} in the path metric on $\widetilde{S} \times \{1\}$.

For the universal cover $\widetilde{B}$ of the thick block B, define:

$$\Pi_{\lambda,B}(x) = \Pi_{Bi}(x), x \in \widetilde{S} \times \{i\}, i = 0, 1$$

Construction of $\Pi_{\lambda,B}$ - Split Block
Definition: Let $y \in Y$ and let μ_q be an electro-ambient representative of an electric geodesic μ_{Gel} in (Y, d_{Gel}). Then $\pi_e(y) = z \in \mu_q$ if the ordered pair $\{d_{Gel}(y, \pi_e(y)), d(y, \pi_e(y))\}$ is minimized at z.

On $\widetilde{S} \times \{0\} \subset \widetilde{B_{tel}}$, let Π_{B0} denote nearest point projection onto λ_{B0}. Here the nearest point projection is taken in the path metric on $\widetilde{S} \times \{0\}$ which is a hyperbolic metric space.

Again, on $\widetilde{S} \times \{1\}$, let Π_{B3} denote nearest point projection onto λ_{B3}. Here the nearest point projection is taken in the path metric on $\widetilde{S} \times \{1\}$ which is a hyperbolic metric space.

For the universal cover $\widetilde{B}$ of the thin block B, define:

$$\Pi_{\lambda,B}(x) = \Pi_{Bi}(x), x \in \widetilde{S} \times \{i\}, i = 0, \cdots, 3$$

$\Pi_{\lambda,B}$ is a retract

Lemma 2.2. *([11] [13]) There exists $C > 0$ such that the following holds: Let $x, y \in \widetilde{S} \times \{0, 1\} \subset \widetilde{B}$ for some block B. Then $d(\Pi_{\lambda,B}(x), \Pi_{\lambda,B}(y)) \leq Cd(x, y)$.*

Construction of B_λ and Π_λ

Given a manifold M of split geometry, we know that M is homeomorphic to $S \times J$ for $J = [0, \infty)$. There exists a sequence I_i of intervals and blocks B_i where the metric on $S \times I_i$ coincides with that on some building block B_i. Denote:
- $B_\mu(B_i) = B_{i\mu}$
- $\Pi_{\mu,B_i} = \Pi_{i\mu}$

Now for a block $B = S \times I$ (thick or split), a natural map Φ_B may be defined taking $\mu = B_\mu(B) \cap \widetilde{S} \times \{0\}$ to a geodesic $B_\mu(B) \cap \widetilde{S} \times \{k\} = \Phi_B(\mu)$ where $k = 1$ or 3 according as B is thick or split. Let the map Φ_{B_i} be denoted as Φ_i for $i \geq 0$.

We start with a reference block B_0 and a reference geodesic segment $\lambda = \lambda_0$ on the 'lower surface' $\widetilde{S} \times \{0\}$. Now inductively define:

- $\lambda_{i+1} = \Phi_i(\lambda_i)$ for $i \geq 0$
- $B_{i\lambda} = B_{\lambda_i}(B_i)$
- $\Pi_{i\lambda} = \Pi_{\lambda_i, B_i}$
- $B_\lambda = \bigcup_i B_{i\lambda}$
- $\Pi_\lambda = \bigcup_i \Pi_{i\lambda}$

Recall that each $\widetilde{S} \times i$ for $i = 0 \cdots m$ is called a **horizontal sheet** of $\widetilde{B}$, where $m = 1$ or 3 according as B is thick or amalgamated. We will restrict our attention to the union of the horizontal sheets $\widetilde{M_H}$ of $\widetilde{M}$ with the metric induced from the graph model.

Clearly, $B_\lambda \subset \widetilde{M_H} \subset \widetilde{M}$, and Π_λ is defined from $\widetilde{M_H}$ to B_λ. Since $\widetilde{M_H}$ is a 'coarse net' in $\widetilde{M}$ (equipped with the *graph model metric*), we will be able to get all the coarse information we need by restricting ourselves to $\widetilde{M_H}$.

By Lemma 2.2, we obtain the fact that each $\Pi_{i\lambda}$ is a retract. Hence assembling all these retracts together, we have the following basic theorem from [11]:

Theorem 2.3. [11] *There exists $C > 0$ such that for any geodesic $\lambda = \lambda_0 \subset \widetilde{S} \times \{0\} \subset \widetilde{B_0}$, the retraction $\Pi_\lambda : \widetilde{M_H} \to B_\lambda$ satisfies:*

Then $d_{Gel}(\Pi_{\lambda,B}(x), \Pi_{\lambda,B}(y)) \leq C d_{Gel}(x,y) + C$.

qi Rays We also have the following from [11].

Lemma 2.4. *There exists $C \geq 0$ such that for $x_i \in \lambda_i$ there exists $x_{i-1} \in \lambda_{i-1}$ with $d_G(x_i, x_{i-1}) \leq C$. Hence, for there exists a C-quasigeodesic ray r such that $r(i) \in \lambda_i \subset B_\lambda$ for $i \geq 0$.*

Further, by construction of split blocks, $d_G(x_i, S_{i-1}) = 1$.

Hence, given $p \in \lambda_i$ the sequence of points $x_n, n \in \mathbb{N} \cup \{0\}$ with $x_i = p$ gives by Lemma 2.4 above, a quasigeodesic in the d_G-metric. Such quasigeodesics shall be referred to as *d_G-quasigeodesic rays*.

3. Laminations

To distinguish between the ending lamination and bi-infinite geodesics whose end-points are identified by ∂i, we make the following definition.

Definition 3.1. *A* **CT leaf** *λ_{CT} is a bi-infinite geodesic whose end-points are identified by ∂i.*
An **EL leaf** *λ_{EL} is a bi-infinite geodesic whose end-points are ideal boundary points of either a leaf of the ending lamination, or a complementary ideal polygon.*

Then to prove the main theorem 1.3 it suffices to show that

- **A** *CT leaf* **is an** *EL leaf***.**

3.1. Leaves of Laminations.

Our first observation is that any semi-infinite geodesic (in the *hyperbolic metric* on $\tilde{S}$) contained in a *CT leaf* in the base surface $\tilde{S}$ $= \tilde{S} \times \{0\} \subset \tilde{B} = \tilde{B} \times \{0\}$ has infinite diameter in the *graph metric* d_G restricted to $\tilde{B}$.

Lemma 3.2. CT leaves have infinite diameter
Let $\lambda_+ (\subset \lambda \subset \tilde{S} \times \{0\} = \tilde{S})$ be a semi-infinite geodesic (in the hyperbolic metric *on $\tilde{S}$) contained in a CT leaf λ. Then $dia_G(\lambda_+)$ is infinite (where dia_G denotes diameter in the graph metric restricted to $\tilde{S}$.*

Proof: Suppose not. Let λ_∞ be the ideal end-point of λ_+. Also let the finite end-point of λ_+ be x, so that $\lambda_+ = [x, \lambda_\infty)$. Then there exist points $x_n \in \lambda_+$ tending to infinity, and a number $K > 0$ such that $d_G(x, x_n) \leq K$.

Also, without loss of generality, assume that $\lambda \subset \widetilde{S} \times \{0\} \widetilde{B_0}$ for the block B_0. Hence there exist split components $\widetilde{B^i} \subset \widetilde{B_0}$, such that $dia_{hyp}(\lambda \cap \widetilde{B^i}) \geq i$ (Here, dia_{hyp} denotes diameter in the hyperbolic metric on $\widetilde{M}$). Acting on $\widetilde{B}$ by elements h_i of the surface group $\pi_1(S)$, we may assume that there exist a sequence of segments $\lambda^i \subset h_i \cdot \lambda$ such that

•1 λ^i is approximately centered about a fixed origin 0, i.e. λ^i pass uniformly close to 0 and end-points of λ^i are at distance $\geq i$ from 0

•2 All the λ^i's lie entirely within a split component $\widetilde{B^i}$

Since $\widetilde{B^i}$ is quasiconvex, it follows that the λ^i's are uniform quasigeodesics in $\widetilde{M}$. Hence, in the limit we obtain a bi-infinite quasigeodesic λ^∞ which is a limit in the Chabauty topology of $h_i \cdot \lambda$'s. Since the set of *CT leaves* are closed in the Chabauty topology, it follows that λ^∞ is a *CT leaf*.

But, this is a contradiction, as we have noted already that λ^∞ is a quasigeodesic. $\Box$

In fact, the proof of Lemma 3.2 shows the following somewhat stronger assertion.

Corollary 3.3. *Given $k \geq 0$, there exists $C \geq 0$ such that if $B = \cup_{0 \leq i \leq k} B_i$ and $\lambda \subset \widetilde{B}$ is a bi-infinite geodesic in the intrinsic metric on $\widetilde{B}$, whose end-points are identified by the Cannon-Thurston map, then for any split component $\widetilde{B^i}$, $dia_{hyp}(\lambda \cap \widetilde{B^i}) \leq C$*

Using Corollary 3.3, we shall now show:

Proposition 3.4. *There exists a function $M(N) \rightarrow \infty$ as $N \rightarrow \infty$ such that the following holds:*
Let λ be a CT leaf. If $a_i, b_i \in \lambda$ be such that $d(a_i, 0) \geq N$, $d(b_i, 0) \geq N$, then $d_G(\overline{a_i b_i}, 0) \geq M(N)$, where d denotes the hyperbolic metric on $\widetilde{M}$.

Proof: Suppose not. Let λ_+ and λ_- denote the ideal end points of λ. Then there exists $C \geq 0$, $a_i \rightarrow \lambda_-$, $b_i \rightarrow \lambda_+$ such that $d_G(\overline{a_i b_i}, 0) \leq C$. That is, there exist $p_i \in \overline{a_i b_i}$ such that $d_G(0, p_i) \leq C$. Due to the existence of a Cannon-Thurston map in the hyperbolic metric (Theorem 1.1), we may assume that $d(0, p_i) \geq i$ (in the hyperbolic metric). Then the hyperbolic geodesic $\overline{0, p_i}$ passes through at most C split blocks for every i. Let $B = \cup_{0 \leq i \leq C} B_i$ and $p_i \rightarrow p_\infty$. Then $\overline{0, p_i} \subset \widetilde{B}$. But since $p_i \in \overline{a_i b_i}$, then the Cannon-Thurston map identifies $\lambda_-, \lambda_+, p_\infty$. See Figure below.



Figure 3: *Cannon-Thurston in the Graph Metric*

Also,

$$\begin{gathered}\overline{0,p_\infty} \subset \overline{0,p_\infty} \cup \overline{0,\lambda_+} \\ \overline{0,p_\infty} \subset \overline{0,p_\infty} \cup \overline{0,\lambda_-}\end{gathered}$$

and at least one of the above two ($\overline{0,p_\infty} \cup \overline{0,\lambda_+} = \overline{p_\infty,\lambda_+}$ say) must pass close to 0. Then $\overline{p_\infty,\lambda_+}$ is a *CT leaf*. But $\overline{0,p_\infty}$ lies in a C-neighborhood of 0 in the graph-metric d_G, contradicting Corollary 3.3 above. □

3.2. **Rays Coarsely Separate Ladders.**

Definition 3.5. *Let X, Y, Z be geodesically complete metric spaces such that $X \subset Y \subset Z$. X is said to* **coarsely separate** *Y into Y_1 and Y_2 if*
(1) $Y_1 \cup Y_2 = Y$
(2) $Y_1 \cap Y_2 = X$
(3) For all $M \geq 0$, there exist $y_1 \in Y_1$ and $y_2 \in Y_2$ such that $d(y_1, Y_2) \geq M$ and $d(y_2, Y_1) \geq M$
(4) There exists $C \geq 0$ such that for all $y_1 \in Y_1$ and $y_2 \in Y_2$ any geodesic in Z joining y_1, y_2 passes through a C-neighborhood of X.

See also [6] for a related notion of coarse separation.

We now fix a quasigeodesic ray r_0 as in Lemma 2.4, and consider a translate $r' = h \cdot r_0$ passing through $z \in \lambda_m \subset B_\lambda$, i.e. $h \cdot r'(m) = z$. Let $\Pi_\lambda \cdot r' = r \subset B_\lambda$. Each $r(i)$ cuts λ_i into two pieces λ_i^- and λ_i^+ with ideal boundary points $\lambda_{i,-\infty}, \lambda_{i,\infty}$ respectively.

We shall show that r coarsely separates B_λ into B_λ^+ and B_λ^-, where

$$\begin{gathered}B_\lambda^+ = \bigcup_i \lambda_i^+ \\ B_\lambda^- = \bigcup_i \lambda_i^-\end{gathered}$$

and λ_i^+ (resp. λ_i^-) is the segment of λ joining $r(i)$ to the ideal end-point $\lambda_{i,-\infty}$ (resp. $\lambda_{i,\infty}$.

We need to repeatedly apply Theorem 2.3 to prove the above assertion.

Given r', we construct two hyperbolic ladders $B_\lambda^{+\prime}$ and $B_\lambda^{-\prime}$, obtained by joining the points $r'(i)$ to the ideal end-points $\lambda_{i,-\infty}, \lambda_{i,\infty}$ of $\lambda_i \subset \widetilde{S} \times \{i\}$. Then $B_\lambda^{+\prime}$ and $B_\lambda^{-\prime}$ are C-quasiconvex by Theorem 2.3. Further, $\Pi_\lambda \cdot r'(i) = r(i)$ by definition of r. Hence,

$$\begin{gathered}\Pi_\lambda(B_\lambda^{-\prime}) = B_\lambda^- \\ \Pi_\lambda(B_\lambda^{+\prime}) = B_\lambda^+\end{gathered}$$

Further,

$$\begin{gathered}B_\lambda^- \cup B_\lambda^+ = B_\lambda \\ B_\lambda^- \cap B_\lambda^+ = r\end{gathered}$$

and there exists a K_0 (independent of r_0, h, λ) such that $B_\lambda^-, B_\lambda^+, B_\lambda, r$ are all K-quasiconvex.

Criterion (3) of Definition 3.5 in this context is given by Lemma 3.2 : *CT leaves have infinite diameter.*

To prove that r separates B_λ into B_λ^-, B_λ^+, we need to show first:

Lemma 3.6. *For all $K_0 \geq 0$, there exists $K_1 \geq 0$ such that if $p \in B_\lambda^-, q \in B_\lambda^+$ with $d_G(p,q) \leq K_0$, then there exists $z \in r$ such that $d_G(p,z) \leq K_1$ and $d_G(q,z) \leq K_1$.*

Proof: Let Π_λ^+ denote the usual (sheetwise)retract (of Theorem 2.3) onto B_λ^+. Then

$$\Pi_\lambda^+(\lambda_i^-) = r(i)$$

and

$$\Pi_\lambda^+(x) = x$$

for all $x \in B_\lambda^+$.

Hence

$$\Pi_\lambda^+(q) = q$$
$$\Pi_\lambda^+(p) = z = r(i)$$

for some $z \in r$ and some i.

Therefore, by Theorem 2.3 again,

$$d_G(q, z) \leq C d_g(p, q) = CK_0$$

.

Choosing $K_1 = CK_0 + K_0$ (and using the triangle inequality for p, q, z) the Lemma follows. □.

We are now in a position to prove:

Theorem 3.7. *r coarsely separates B_λ into B_λ^-, B_λ^+.*

Proof: We have already shown

$$B_\lambda^- \cup B_\lambda^+ = B_\lambda$$
$$B_\lambda^- \cap B_\lambda^+ = r$$

and there exists a K_0 (independent of r_0, h, λ) such that $B_\lambda^-, B_\lambda^+, B_\lambda, r$ are all K-quasiconvex.

Criterion (3) of Definition 3.5 is given by Lemma 3.2.

Finally given $u \in B_\lambda^-$ and $v \in B_\lambda^+$, let $\overline{uv}$ be the geodesic in $(\widetilde{M}, d_G)$ joining u, v. Then $\Pi_\lambda(\overline{uv})$ is a "dotted quasigeodesic" $u = p_0, p_1, \cdots p_n = v$, where $d_G(p_i, p_{i+1}) \leq C$ (and C is obtained from Theorem 2.3). Further, $p_0 \in B_\lambda^-, p_n \in B_\lambda^+$ and $p_i \in B_\lambda$ for all i. Therefore there exists m such that $p_m \in B_\lambda^-, p_{m+1} \in B_\lambda^+$, with $d_G(p_m, p_{m+1}) \leq C$. Hence, by Lemma 3.6, there exists $K_1 \geq 0$ such that there exists $z \in r$ with $d_G(p_m, z) \leq K_1$ and $d_G(p_{m+1}, z) \leq K_1$.

Finally, by Definition-Theorem Split Geometry 2.1, $(\widetilde{M}, d_G)$ is hyperbolic, and therefore the "dotted quasigeodesic" $u = p_0, p_1, \cdots p_n = v$ lies in a uniformly bounded neighborhood of the geodesic $\overline{uv}$. That is, there exists $C_1 \geq 0$ such that for all $u \in B_\lambda^-$ and $v \in B_\lambda^+$, the geodesic $\overline{uv}$ in $(\widetilde{M}, d_G)$ joining u, v passes through a C_1-neighborhood of r. This proves (4) in Definition 3.5 and hence we conclude that r coarsely B_λ into B_λ^-, B_λ^+. □

We shall have need for the following Proposition, whose proof is along the lines of Theorem 3.7 above.

Proposition 3.8. *Let μ, λ be two bi-infinite geodesics on $\widetilde{S}$ such that $\mu \cap \lambda \neq \emptyset$. Then $B_\lambda \cap B_\mu = r$ is a quasigeodesic ray coarsely separating both B_λ and B_μ.*

Next, by Proposition 3.4, we find that if $a_i, b_i \in \lambda = \lambda_0$ such that a_i, b_i converge to ideal points $\lambda_{0,-\infty}, \lambda_{0,\infty}$ (denoted $\lambda_{-\infty}, \lambda_\infty$ for convenience), then $\Pi_\lambda(\overline{a_i b_i})$ leaves large balls about 0. Also, by Theorem 3.7 above, r coarsely separates B_λ. Hence $\Pi_\lambda(\overline{a_i b_i})$ passes close to $r(n_i)$ for some $n_i \in \mathbb{N}$, where $n_i \to \infty$ as $i \to \infty$. Hence, we conclude that any such r converges on $\partial\widetilde{M}$ to the same point as $\partial(\lambda_{-\infty}) = \partial(\lambda_\infty)$. In particular any two quasigeodesic rays lying on B_λ are asymptotic. We thus have:

Proposition 3.9. Asymptotic Quasigeodesic Rays
Given $K \geq 1$ there exists α such that if λ is a CT-leaf then there exists $z \in \partial\widetilde{M}$ satisfying the following:
If r_1 and r_2 are K-quasi-isometric sections contained in B_λ then there exists $N \in \mathbb{N}$ such that
1) $r_j(n) \rightarrow z = \partial i(\lambda_{-\infty}) = \partial i(\lambda_\infty)$ as $n \rightarrow \infty$, for $j = 1, 2$.
2) $d_G(r_1(n), r_2(n)) \leq \alpha$ for all $n \geq N$

3.3. **Main Theorem.** We are now in a position to prove the main Theorem of this paper.

Main Theorem 1.3 *Let $\partial i(a) = \partial i(b)$ for $a, b \in S^1_\infty$ be two distinct points that are identified by the Cannon-Thurston map corresponding to a simply degenerate closed surface group (without accidental parabolics). Then a, b are either ideal end-points of a leaf of the ending lamination (in the sense of Thurston), or ideal boundary points of a complementary ideal polygon. Conversely, if a, b are either ideal end-points of a leaf of a lamination, or ideal boundary points of a complementary ideal polygon, then $\partial i(a) = \partial i(b)$.*

Proof of Main Theorem 1.3: The converse direction has been shown in Proposition 1.4.
Next, let $\partial i(a) = \partial i(b)$ for $a, b \in S^1_\infty$. Then $(a, b) = \lambda \subset \widetilde{S_0} \subset \widetilde{M}$ is a CT-leaf.

Suppose λ and μ are intersecting CT leaves, i.e. $\partial i(\lambda_{-\infty}) = \partial i(\lambda_\infty)$ and $\partial i(\mu_{-\infty}) = \partial i(\mu_\infty)$.

As before, let λ_i and μ_i be intersections of the ladders B_λ and B_μ with the horizontal sheets. Let $r(i) = \lambda_i \cap \mu_i$ be a quasigeodesic ray as per Proposition 3.8. Hence by Proposition 3.9, r converges to a point z on $\partial\widetilde{M}$ such that $z = \partial i(\lambda_{-\infty}) = \partial i(\lambda_\infty) = \partial i(\mu_{-\infty}) = \partial i(\mu_\infty)$. Hence we have shown that the Cannon-Thurston map identies the endpoints of any two intersecting CT leaves λ and μ.

If possible, suppose that the CT leaf λ is not an EL-leaf. Then λ intersects the ending lamination transversely (since the ending lamination is a filling lamination without any closed leaves) and there exist EL-leaves μ for which $\lambda \cap \mu \neq \emptyset$. By Proposition 1.4, each such μ is a CT-leaf. Hence, by the previous paragraph, the Cannon- Thurston map identies the end points of λ with the endpoints of each such EL-leaf μ for which $\lambda \cap \mu \neq \emptyset$.

Since λ is not an EL-leaf, it contains a non-trivial geodesic subsegment I transverse to the ending lamination.

Let Λ denote the union of all lifts of leaves of the ending lamination to the universal cover $\widetilde{S}$. Let $\widetilde{I}$ denote the union of all lifts of I to $\widetilde{S}$. Define two leaves of $\widetilde{\Lambda}$ to be equivalent if both of them intersect a single element of $\widetilde{I}$. Then by Lemma 1.9 the limit set $\mathcal{L}$ of any connected component of the transitive closure of this relation contains a pole.

All the points of $\mathcal{L}$ are identified to a point by the existence of Cannon-Thurston maps (Theorem 1.1) and by Proposition 1.4. Hence, in particular the Cannon-Thurston map identifies a pole to a point. This is a contradiction as a pole forms the end-points of a quasigeodesic in $\widetilde{M}$.

This contradiction shows that λ must be an EL-leaf. $\square$

Totally Degenerate Groups We have so far assumed, for ease of exposition, that we are dealing with simply degenerate groups. A few words are in order as to the necessary modifications to be made for the totally degenerate case. All that we need to do to generalize to this situation is fix a base reference surface. This is the only place where we used the boundary of the convex core as a preferred base reference surface, but any fixed pleated surface would do as well. The same works for 3-manifolds with incompressible boundary. Here, we fix a reference base surface for each end, and use the existence of Cannon-Thurston maps, ladders B_λ and rays from [11].

References

[1] F. Bonahon. Bouts de varietes hyperboliques de dimension 3. *Ann. Math. vol.124*, pages 71–158, 1986.

[2] R. D. Canary, D. B. A. Epstein, and P. Green. Notes on Notes of Thurston. *in Analytical and Geometric Aspects of Hyperbolic Spaces*, pages 3–92, 1987.

[3] J. Cannon and W. P. Thurston. Group Invariant Peano Curves. *preprint, Princeton*, 1985.

[4] A. Casson and S. Bleiler. Automorphisms of Surfaces after Nielsen and Thurston. *London Math. Soc. Student Texts, Cambridge*, 1987.

[5] B. Farb. Relatively hyperbolic groups. *Geom. Funct. Anal. 8*, pages 810–840, 1998.

[6] B. Farb and R. E. Schwartz. Quasi-isometric Rigidity of Hilbert Modular Groups. *Jour. Diff. Geom. 44(3)*, page 435478, 1996.

[7] E. Ghys and P. de la Harpe(eds.). Sur les groupes hyperboliques d'apres Mikhael Gromov. *Progress in Math. vol 83, Birkhauser, Boston Ma.*, 1990.

[8] E. Klarreich. Semiconjugacies between Kleinian group actions on the Riemann sphere. *Amer. J. Math 121*, pages 1031–1078, 1999.

[9] Y. N. Minsky. On Rigidity, Limit Sets, and End Invariants of Hyperbolic 3-Manifolds. *J. A.M.S., vol.7*, pages 539–588, 1994.

[10] Mahan Mitra. Ending Laminations for Hyperbolic Group Extensions. *GAFA vol.7 No. 2*, pages 379–402, 1997.

[11] Mahan Mj. Cannon-Thurston Maps for Surface Groups I: Amalgamation Geometry and Split Geometry. *preprint, arXiv:math.GT/0512539*, 2005.

[12] Mahan Mj. Cannon-Thurston Maps, i-bounded Geometry and a Theorem of McMullen. *preprint, arXiv:math.GT/0511041*, 2005.

[13] Mahan Mj. Cannon-Thurston Maps for Surface Groups II: Split Geometry and the Minsky Model. *preprint, arXiv:math.GT/*, 2006.

[14] R. Penner and J. Harer. Combinatorics of train tracks. *Ann. Math. Studies 125, Princeton University Press*, 1992.

[15] Peter B. Shalen. Dendrology and its applications. *Group Theory from a Geometrical Viewpoint (E. Ghys, A. Haefliger, A. Verjovsky eds.)*, page 543616, 1991.

[16] W. P. Thurston. The Geometry and Topology of 3-Manifolds. *Princeton University Notes*, 1980.

RKM Vivekananda University, Belur Math, WB-711202, India
E-mail address: `mahanDOTmjATgmailDOTcom`